arXiv:1205.0716v1 [math.DG] 3 May 2012

Jet Riemann-Hamilton geometrization for the conformal deformed Berwald-Moór quartic metric depending on momenta

Alexandru Oană and Mircea Neagu

Abstract

In this paper we expose on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ the distinguished (d-) Riemannian geometry (in the sense of d-connection, d-torsions, d-curvatures and some gravitational-like and electromagnetic-like geometrical models) for the (t, x)-conformal deformed Berwald-Moór Hamiltonian metric of order four.

Mathematics Subject Classification (2010): 70S05, 53C07, 53C80.

Key words and phrases: (t, x)-conformal deformation of quartic Berwald-Moór Hamiltonian metric, canonical nonlinear connection, Cartan canonical linear connection, d-torsions and d-curvatures, geometrical Einstein-like equations.

1 Introduction

The geometric-physical Lagrangian or Hamiltonian Berwald-Moór structure ([4], [7]) was intensively investigated by P.K. Rashevski ([13]) and further fundamented and developed by D.G. Pavlov, G.I. Garas'ko and S.S. Kokarev ([11], [5], [12]). In their works, the preceding Russian scientists emphasize the importance in the theory of space-time structure, gravitation and electromagnetism of the geometry produced by the classical Berwald-Moór Lagrangian metric

$$F : TM \to \mathbb{R}, \qquad F(y) = \sqrt[n]{y^1 y^2 ... y^n}, \qquad n \geq 2,$$

or by the corresponding Berwald-Moór Hamiltonian metric

$$H : T^*M \to \mathbb{R}, \qquad H(p) = \sqrt[n]{p_1 p_2 ... p_n}.$$

In such a perspective, according to the recent geometric-physical ideas proposed by Garas'ko ([5]), we consider that a distinguished Riemannian geometry (in the sense of d-connection, d-torsions, d-curvatures and some gravitational-like and electromagnetic-like geometrical models) for the conformal deformations of the jet Berwald-Moór Hamiltonian metric of order four is required. Note that a similar geometric-physical study for the (t, x)-conformal deformations of the jet Berwald-Moór Lagrangian metric of order four is now completely developed in the paper [8]. Also, few elements of distinguished Hamiltonian geometry produced by the cotangent quartic Berwald-Moór metric depending of momenta are already presented in the paper [1].

In such a geometrical and physical context, this paper investigates on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ the Riemann-Hamilton distinguished geometry (together with a theoretical-geometric field-like theory) for the (t,x)-*conformal deformed Berwald-Moór Hamiltonian metric of order four*[1]

$$H(t,x,p) = 2e^{-\sigma(x)}\sqrt{h_{11}(t)}\left[p_1^1 p_2^1 p_3^1 p_4^1\right]^{1/4}, \tag{1.1}$$

where $\sigma(x)$ is a smooth non-constant function on M^4, $h_{11}(t)$ is a Riemannian metric on $\mathbb{R}$, and $(t,x,p) = (t, x^1, x^2, x^3, x^4, p_1^1, p_2^1, p_3^1, p_4^1)$ are the coordinates of the momentum phase space $J^{1*}(\mathbb{R}, M^4)$; these transform by the rules (the Einstein convention of summation is assumed everywhere):

$$\widetilde{t} = \widetilde{t}(t), \quad \widetilde{x}^i = \widetilde{x}^i(x^j), \quad \widetilde{p}_i^1 = \frac{\partial x^j}{\partial \widetilde{x}^i}\frac{d\widetilde{t}}{dt}p_j^1, \tag{1.2}$$

where $i,j = \overline{1,4}$, rank $(\partial\widetilde{x}^i/\partial x^j) = 4$ and $d\widetilde{t}/dt \neq 0$. It is important to note that, based on the geometrical ideas promoted by Miron, Hrimiuc, Shimada and Sabău in the classical Hamiltonian geometry of cotangent bundles ([6]), together with those used by Atanasiu, Neagu and Oană in the geometry of dual 1-jet spaces, the differential geometry (in the sense of d-connections, d-torsions, d-curvatures, abstract gravitational-like and electromagnetic-like geometrical theories) produced by a jet Hamiltonian function $H : J^{1*}(\mathbb{R}, M^n) \to \mathbb{R}$ is now completely done in the papers [2], [3], [9] and [10]. In what follows, we apply the general geometrical results from [9] and [10] to the square of Hamiltonian metric (1.1), which is given by $(n = 4)$

$$\overset{*}{H}(t,x,p) = H^2(t,x,p) = 4e^{-2\sigma(x)}h_{11}(t)\left[p_1^1 p_2^1 p_3^1 p_4^1\right]^{1/2}. \tag{1.3}$$

Remark 1.1 *The momentum Hamiltonian metric (1.3) is exactly the natural Hamiltonian attached to the square of the jet Berwald-Moór Lagrangian metric, which has the expression*

$$\overset{*}{L}(t,x,y) = e^{2\sigma(x)}h^{11}(t)\left[y_1^1 y_1^2 y_1^3 y_1^4\right]^{1/2}. \tag{1.4}$$

In other words, we have

$$\overset{*}{H}(t,x,p) = p_r^1 y_1^r - \overset{*}{L}(t,x,y),$$

where $p_i^1 = \partial\overset{*}{L}/\partial y_1^i$. *Note that the jet Lagrangian metric (1.4) is even the square of the conformal deformed jet quartic Berwald-Moór Finslerian metric*

$$F(t,x,y) = e^{\sigma(x)}\sqrt{h^{11}(t)}\left[y_1^1 y_1^2 y_1^3 y_1^4\right]^{1/4}.$$

In other words, we have $\overset{*}{L} = F^2$.

---

[1] We assume that we have $p_1^1 p_2^1 p_3^1 p_4^1 > 0$. This is one domain where we can p-differentiate the Hamiltonian function $H(t,x,p)$.

2 The canonical nonlinear connection

Using the notation $\mathcal{P}^{1111} := p_1^1 p_2^1 p_3^1 p_4^1$ and taking into account that we have

$$\frac{\partial \mathcal{P}^{1111}}{\partial p_i^1} = \frac{\mathcal{P}^{1111}}{p_i^1},$$

then the *fundamental metrical d-tensor* produced by the metric (1.3) is given by the formula (no sum by i or j)

$$\overset{*}{g}{}^{ij}(t,x,p) = \frac{h^{11}(t)}{2}\frac{\partial^2 \overset{*}{H}}{\partial p_i^1 \partial p_j^1} = \frac{e^{-2\sigma(x)}\left(1-2\delta^{ij}\right)}{2}\frac{\left[\mathcal{P}^{1111}\right]^{1/2}}{p_i^1 p_j^1}. \tag{2.1}$$

Moreover, the matrix $\overset{*}{g} = (\overset{*}{g}{}^{ij})$ admits the inverse $\overset{*}{g}{}^{-1} = (\overset{*}{g}_{jk})$, whose entries are given by

$$\overset{*}{g}_{jk} = \frac{e^{2\sigma(x)}\left(1-2\delta_{jk}\right)\left[\mathcal{P}^{1111}\right]^{-1/2}}{2} p_j^1 p_k^1 \text{ (no sum by } j \text{ or } k). \tag{2.2}$$

Let us consider the Christoffel symbol of the Riemannian metric $h_{11}(t)$ on $\mathbb{R}$, which is given by

$$\mathrm{K}_{11}^1 = \frac{h^{11}}{2}\frac{dh_{11}}{dt},$$

where $h^{11} = 1/h_{11} > 0$. Then, using the notation $\sigma_i := \partial\sigma/\partial x^i$, we find the following geometrical result:

Proposition 2.1 *For the (t,x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3), the* ***canonical nonlinear connection*** *on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ has the components*

$$N = \left(\underset{1}{N}{}_{(i)1}^{(1)} = \mathrm{K}_{11}^1 p_i^1,\ \underset{2}{N}{}_{(i)j}^{(1)} = -4\sigma_i p_i^1 \delta_{ij}\right). \tag{2.3}$$

Proof. The canonical nonlinear connection produced by $\overset{*}{H}$ on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ has the following components (see [2]): $\underset{1}{N}{}_{(i)1}^{(1)} \overset{def}{=} \mathrm{K}_{11}^1 p_i^1$ and

$$\underset{2}{N}{}_{(i)j}^{(1)} = \frac{h^{11}}{4}\left[\frac{\partial \overset{*}{g}_{ij}}{\partial x^k}\frac{\partial \overset{*}{H}}{\partial p_k^1} - \frac{\partial \overset{*}{g}_{ij}}{\partial p_k^1}\frac{\partial \overset{*}{H}}{\partial x^k} + \overset{*}{g}_{ik}\frac{\partial^2 \overset{*}{H}}{\partial x^j \partial p_k^1} + \overset{*}{g}_{jk}\frac{\partial^2 \overset{*}{H}}{\partial x^i \partial p_k^1}\right].$$

Now, by a direct calculation, we obtain (2.3). ∎

3 The Cartan canonical N-linear connection. Its d-torsions and d-curvatures

The nonlinear connection (2.3) produces the dual *adapted bases* of d-vector fields (no sum by i)

$$\left\{\frac{\delta}{\delta t} = \frac{\partial}{\partial t} - \mathrm{K}_{11}^1 p_r^1 \frac{\partial}{\partial p_r^1}\ ;\ \frac{\delta}{\delta x^i} = \frac{\partial}{\partial x^i} + 4\sigma_i p_i^1 \frac{\partial}{\partial p_i^1}\ ;\ \frac{\partial}{\partial p_i^1}\right\} \subset \mathcal{X}(E^*) \tag{3.1}$$

and d-covector fields (no sum by i)

$$\left\{dt\ ;\ dx^i\ ;\ \delta p_i^1 = dp_i^1 + \varkappa_{11}^1 p_i^1 dt - 4\sigma_i p_i^1 dx^i\right\} \subset \mathcal{X}^*(E^*), \tag{3.2}$$

where $E^* = J^{1*}(\mathbb{R}, M^4)$. The naturalness of the geometrical adapted bases (3.1) and (3.2) is coming from the fact that, via a transformation of coordinates (1.2), their elements transform as the classical tensors. Therefore, the description of all subsequent geometrical objects on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ (e.g., the Cartan canonical linear connection, its torsion and curvature etc.) will be done in local adapted components. Consequently, by direct computations, we obtain the following geometrical result:

Proposition 3.1 *The Cartan canonical N-linear connection produced by the (t,x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) has the following adapted local components (no sum by i, j or k):*

$$C\Gamma(N) = \left(\varkappa_{11}^1,\ A_{j1}^i = 0,\ H_{jk}^i = 4\delta_j^i\delta_k^i\sigma_i,\ C_{i(1)}^{j(k)} = \mathsf{C}_i^{jk}\cdot\frac{p_i^1}{p_j^1 p_k^1}\right), \tag{3.3}$$

where

$$\begin{aligned}\mathsf{C}_i^{jk} &= \frac{1 - 2\delta^{jk} - 2\delta_i^j - 2\delta_i^k + 8\delta_i^j\delta_i^k}{8} = \\ &= \begin{cases} \dfrac{1}{8}, & i \neq j \neq k \neq i \\ -\dfrac{1}{8}, & i = j \neq k \ \text{or}\ i = k \neq j\ \text{or}\ j = k \neq i \\ \dfrac{3}{8}, & i = j = k. \end{cases}\end{aligned}$$

Proof. The adapted components of the Cartan canonical connection are given by the formulas (see [9]),

$$A_{j1}^i \overset{def}{=} \frac{\overset{*}{g}{}^{il}}{2}\frac{\delta \overset{*}{g}_{lj}}{\delta t} = 0, \quad H_{jk}^i \overset{def}{=} \frac{\overset{*}{g}{}^{ir}}{2}\left(\frac{\delta\overset{*}{g}_{jr}}{\delta x^k} + \frac{\delta\overset{*}{g}_{kr}}{\delta x^j} - \frac{\delta\overset{*}{g}_{jk}}{\delta x^r}\right) = 4\delta_j^i\delta_k^i\sigma_i,$$

$$C_{i(1)}^{j(k)} \overset{def}{=} -\frac{\overset{*}{g}_{ir}}{2}\left(\frac{\partial\overset{*}{g}{}^{jr}}{\partial p_k^1} + \frac{\partial\overset{*}{g}{}^{kr}}{\partial p_j^1} - \frac{\partial\overset{*}{g}{}^{jk}}{\partial p_r^1}\right) = -\frac{\overset{*}{g}_{ir}}{2}\frac{\partial\overset{*}{g}{}^{jr}}{\partial p_k^1}.$$

Using the derivative operators (3.1), the direct calculations lead us to the required results. ∎

Remark 3.2 *It is important to note that the vertical d-tensor $C_{j(k)}^{i(1)}$ also has the properties (sum by m):*

$$C_{i(1)}^{j(k)} = C_{i(1)}^{k(j)}, \quad C_{i(1)}^{j(m)}p_m^1 = 0, \quad C_{m(1)}^{j(m)} = 0, \quad C_{i(1)|m}^{j(m)} = 0, \tag{3.4}$$

where

$$C_{i(1)|j}^{l(k)} \overset{def}{=} \frac{\delta C_{i(1)}^{l(k)}}{\delta x^j} + C_{i(1)}^{r(k)}H_{rj}^l - C_{r(1)}^{l(k)}H_{ij}^r + C_{i(1)}^{l(r)}H_{rj}^k.$$

Proposition 3.3 *The Cartan canonical connection of the (t,x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) has **two** effective local torsion d-tensors:*

$$R^{(1)}_{(r)ij} = -4\sigma_{ij}\left(p^1_i\delta_{ir} - p^1_j\delta_{jr}\right), \quad P^{r(j)}_{i(1)} = \mathsf{C}^{rj}_{i} \cdot \frac{p^1_i}{p^1_r p^1_j},$$

where $\sigma_{ij} := \dfrac{\partial^2\sigma}{\partial x^i \partial x^j}$.

Proof. A Cartan canonical connection on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ generally has *six* effective local d-tensors of torsion (for more details, see [9]). For the particular Cartan canonical connection (3.3) these reduce only to *two* (the other four are zero):

$$R^{(1)}_{(r)ij} \overset{def}{=} \frac{\delta \underset{2}{N}{}^{(1)}_{(r)i}}{\delta x^j} - \frac{\delta \underset{2}{N}{}^{(1)}_{(r)j}}{\delta x^i}, \quad P^{r(j)}_{i(1)} \overset{def}{=} C^{r(j)}_{i(1)}.$$

■

Proposition 3.4 *The Cartan canonical connection of the (t,x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) has **three** effective local curvature d-tensors:*

$$R^l_{ijk} = \frac{\partial H^l_{ij}}{\partial x^k} - \frac{\partial H^l_{ik}}{\partial x^j} + H^r_{ij}H^l_{rk} - H^r_{ik}H^l_{rj} + C^{l(r)}_{i(1)}R^{(1)}_{(r)jk}, \quad P^{l\ (k)}_{ij(1)} = -C^{l(k)}_{i(1)|j},$$

$$S^{l(j)(k)}_{i(1)(1)} \overset{def}{=} \frac{\partial C^{l(j)}_{i(1)}}{\partial p^1_k} - \frac{\partial C^{l(k)}_{i(1)}}{\partial p^1_j} + C^{r(j)}_{i(1)}C^{l(k)}_{r(1)} - C^{r(k)}_{i(1)}C^{l(j)}_{r(1)}.$$

Proof. A Cartan canonical connection on the dual 1-jet space $J^{1*}(\mathbb{R}, M^4)$ generally has *five* effective local d-tensors of curvature (for all details, see [9]). For the particular Cartan canonical connection (3.3) these reduce only to *three* (the other two are zero). These are $S^{l(j)(k)}_{i(1)(1)}$ and

$$R^l_{ijk} \overset{def}{=} \frac{\delta H^l_{ij}}{\delta x^k} - \frac{\delta H^l_{ik}}{\delta x^j} + H^r_{ij}H^l_{rk} - H^r_{ik}H^l_{rj} + C^{l(r)}_{i(1)}R^{(1)}_{(r)jk},$$

$$P^{l\ (k)}_{ij(1)} \overset{def}{=} \frac{\partial H^l_{ij}}{\partial p^1_k} - C^{l(k)}_{i(1)|j} + C^{l(r)}_{i(1)}P^{(1)\ (k)}_{(r)j(1)} = -C^{l(k)}_{i(1)|j},$$

where

$$P^{(1)\ (k)}_{(r)j(1)} = \frac{\partial \underset{2}{N}{}^{(1)}_{(r)j}}{\partial p^1_k} + H^k_{rj} = 0.$$

■

4 From (t, x)-conformal deformations of the quartic Berwald-Moór Hamiltonian metric to field-like geometrical models

4.1 Momentum gravitational-like geometrical model

The (t, x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) produces on the momentum phase space $J^{1*}(\mathbb{R}, M^4)$ the adapted metrical d-tensor (sum by i and j)

$$\mathbb{G} = h_{11}dt \otimes dt + \overset{*}{g}_{ij}dx^i \otimes dx^j + h_{11}\overset{*}{g}^{ij}\delta p_i^1 \otimes \delta p_j^1, \tag{4.1}$$

where $\overset{*}{g}_{jk}$ and $\overset{*}{g}^{ij}$ are given by (2.2) and (2.1), and we have (no sum by i) $\delta p_i^1 = dp_i^1 + \mathrm{K}_{11}^1 p_i^1 dt - 4\sigma_i p_i^1 dx^i$. We believe that, from a physical point of view, the metrical d-tensor (4.1) may be regarded as a *"gravitational potential depending on momenta"*. In our abstract geometric-physical approach, one postulates that the momentum gravitational potential $\mathbb{G}$ is governed by the *geometrical Einstein equations*

$$\text{Ric } (C\Gamma(N)) - \frac{\text{Sc } (C\Gamma(N))}{2}\mathbb{G} = \mathcal{K}\mathbb{T}, \tag{4.2}$$

where

- Ric $(C\Gamma(N))$ is the *Ricci d-tensor* associated to the Cartan canonical linear connection (3.3); the Cartan canonical linear connection plays in our geometric-physical theory the same role as the Levi-Civita connection in the classical Riemannian theory of gravity;
- Sc $(C\Gamma(N))$ is the *scalar curvature*;
- $\mathcal{K}$ is the *Einstein constant* and $\mathbb{T}$ is an intrinsic *momentum stress-energy d-tensor of matter.*

Therefore, using the adapted basis of vector fields (3.1), we can locally describe the global geometrical Einstein equations (4.2). Consequently, some direct computations lead to:

Lemma 4.1 *The Ricci tensor of the Cartan canonical connection of the (t, x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) has the following* ***two*** *effective local Ricci d-tensors (no sum by i, j, k or l):*

$$R_{ij} = \begin{cases} -2\sigma_{ij} - \dfrac{p_i^1}{p_k^1}\sigma_{jk} - \dfrac{p_i^1}{p_l^1}\sigma_{jl}, & i \neq j, \ \ \{i, j, k, l\} = \{1, 2, 3, 4\} \\ 0, & i = j, \end{cases} \tag{4.3}$$

$$R_{(1)(1)}^{(i)(j)} = -S_{(1)(1)}^{(i)(j)} = \frac{4\delta^{ij} - 1}{8}\frac{1}{p_i^1 p_j^1}.$$

Proof. Generally, the Ricci tensor of a Cartan canonical connection $C\Gamma(N)$ (produced by an arbitrary momentum Hamiltonian function) is determined by *six* effective local Ricci d-tensors (for more details, see [10]). For the particular Cartan canonical connection (3.3) these reduce only to *two* (the other four are zero), where (sum by r and m):

$$\begin{aligned} R_{ij} &\overset{def}{=} R^m_{ijm} = \frac{\partial H^m_{ij}}{\partial x^m} - \frac{\partial H^m_{im}}{\partial x^j} + H^r_{ij}H^m_{rm} - H^r_{im}H^m_{rj} + C^{m(r)}_{i(1)}R^{(1)}_{(r)jm}, \\ S^{(i)(j)}_{(1)(1)} &\overset{def}{=} S^{i(j)(m)}_{m(1)(1)} = \frac{\partial C^{i(j)}_{m(1)}}{\partial p^1_m} - \frac{\partial C^{i(m)}_{m(1)}}{\partial p^1_j} + C^{r(j)}_{m(1)}C^{i(m)}_{r(1)} - C^{r(m)}_{m(1)}C^{i(j)}_{r(1)} = \\ &= \frac{\partial C^{i(j)}_{m(1)}}{\partial p^1_m} + C^{r(j)}_{m(1)}C^{i(m)}_{r(1)}. \end{aligned}$$

■

Lemma 4.2 *The scalar curvature of the Cartan canonical connection of the (t,x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) has the value*

$$\text{Sc }(C\Gamma(N)) = -4e^{-2\sigma}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11} - \frac{3}{2}h^{11}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2},$$

where

$$\Sigma_{11} = \frac{\sigma_{12}}{p^1_1p^1_2} + \frac{\sigma_{13}}{p^1_1p^1_3} + \frac{\sigma_{14}}{p^1_1p^1_4} + \frac{\sigma_{23}}{p^1_2p^1_3} + \frac{\sigma_{24}}{p^1_2p^1_4} + \frac{\sigma_{34}}{p^1_3p^1_4}.$$

Proof. The scalar curvature of the Cartan canonical connection (3.3) is given by the general formula

$$\text{Sc }(C\Gamma(N)) = \overset{*}{g}{}^{ij}R_{ij} - h^{11}\overset{*}{g}_{ij}S^{(i)(j)}_{(1)(1)}.$$

■

The local description in the adapted basis of vector fields (3.1) of the global geometrical Einstein equations (4.2) leads us to

Proposition 4.3 *The **geometrical Einstein-like equations** produced by the (t,x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) are locally described by*

$$\begin{cases} 2e^{-2\sigma}h_{11}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11} + \dfrac{3}{4}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2} = \mathcal{K}\mathbb{T}_{11}, \\ R_{ij} + \left\{2e^{-2\sigma}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11} + \dfrac{3}{4}h^{11}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2}\right\}\overset{*}{g}_{ij} = \mathcal{K}\mathbb{T}_{ij}, \\ -S^{(i)(j)}_{(1)(1)} + \left\{ 2e^{-2\sigma}h_{11}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11} + \right. \\ \qquad\qquad \left. + \dfrac{3}{4}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2}\right\}\overset{*}{g}{}^{ij} = \mathcal{K}\mathbb{T}^{(i)(j)}_{(1)(1)}, \\ 0 = \mathbb{T}_{1i}, \quad 0 = \mathbb{T}_{i1}, \quad 0 = \mathbb{T}^{(i)}_{(1)1}, \\ 0 = \mathbb{T}_{1(1)}^{\ \ (i)}, \quad 0 = \mathbb{T}_{i(1)}^{\ \ (j)}, \quad 0 = \mathbb{T}^{(i)}_{(1)j}. \end{cases} \tag{4.4}$$

Corollary 4.4 *The momentum stress-energy d-tensor of matter $\mathcal{T}$ satisfies the following* ***geometrical conservation-like laws*** *(sum by m):*

$$\begin{cases} \mathbb{T}^1_{1/1} + \mathbb{T}^m_{1|m} + \mathbb{T}^{(1)}_{(m)1}\big|^{(m)}_{(1)} = 0 \\ \\ \mathbb{T}^1_{i/1} + \mathbb{T}^m_{i|m} + \mathbb{T}^{(1)}_{(m)i}\big|^{(m)}_{(1)} = \mathsf{E}^m_{i|m} \\ \\ \mathbb{T}^{1(i)}_{\ \ (1)/1} + \mathbb{T}^{m(i)}_{\ \ \ (1)|m} + \mathbb{T}^{(1)(i)}_{(m)(1)}\big|^{(m)}_{(1)} = \dfrac{e^{-2\sigma}\left[\mathcal{P}^{1111}\right]^{1/2}}{\mathcal{K}} \cdot \left[\dfrac{\Sigma_{11}}{p^1_i} + 2\dfrac{\partial \Sigma_{11}}{\partial p^1_i}\right], \end{cases}$$

where (sum by r):

$$\mathbb{T}^1_1 \overset{def}{=} h^{11}\mathbb{T}_{11} = \mathcal{K}^{-1}\left\{2e^{-2\sigma}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11} + \frac{3}{4}h^{11}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2}\right\},$$

$$\mathbb{T}^m_1 \overset{def}{=} \overset{*}{g}{}^{mr}\mathbb{T}_{r1} = 0, \quad \mathbb{T}^{(1)}_{(m)1} \overset{def}{=} h^{11}\overset{*}{g}_{mr}\mathbb{T}^{(r)}_{(1)1} = 0, \quad \mathbb{T}^1_i \overset{def}{=} h^{11}\mathbb{T}_{1i} = 0,$$

$$\mathbb{T}^m_i \overset{def}{=} \overset{*}{g}{}^{mr}\mathbb{T}_{ri} = \mathsf{E}^m_i := \mathcal{K}^{-1}\cdot\left[\overset{*}{g}{}^{mr}R_{ri} + \delta^m_i\left\{2e^{-2\sigma}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11} + \right.\right.$$

$$\left.\left. + \frac{3}{4}h^{11}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2}\right\}\right],$$

$$\mathbb{T}^{(1)}_{(m)i} \overset{def}{=} h^{11}\overset{*}{g}_{mr}\mathbb{T}^{(r)}_{(1)i} = 0, \ \ \mathbb{T}^{1(i)}_{\ \ (1)} \overset{def}{=} h^{11}\mathbb{T}_{1(1)}^{\ \ (i)} = 0, \ \ \mathbb{T}^{m(i)}_{\ \ \ (1)} \overset{def}{=} \overset{*}{g}{}^{mr}\mathbb{T}_{r(1)}^{\ \ (i)} = 0,$$

$$\mathbb{T}^{(1)(i)}_{(m)(1)} \overset{def}{=} h^{11}\overset{*}{g}_{mr}\mathbb{T}^{(r)(i)}_{(1)(1)} = \frac{h^{11}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2}}{8\mathcal{K}}\frac{p_m}{p_i} +$$

$$+\delta^i_m \cdot \left[\frac{h^{11}e^{2\sigma}\left[\mathcal{P}^{1111}\right]^{-1/2}}{4\mathcal{K}} + \frac{2e^{-2\sigma}\left[\mathcal{P}^{1111}\right]^{1/2}\Sigma_{11}}{\mathcal{K}}\right],$$

and we also have (summation by m and r, but no sum by i)

$$\mathbb{T}^1_{1/1} \overset{def}{=} \frac{\delta \mathbb{T}^1_1}{\delta t} + \mathbb{T}^1_1 \mathrm{K}^1_{11} - \mathbb{T}^1_1 \mathrm{K}^1_{11} = \frac{\delta \mathbb{T}^1_1}{\delta t}, \quad \mathbb{T}^m_{1|m} \overset{def}{=} \frac{\delta \mathbb{T}^m_1}{\delta x^m} + \mathbb{T}^r_1 H^m_{rm},$$

$$\mathbb{T}^{(1)}_{(m)1}\big|^{(m)}_{(1)} \overset{def}{=} \frac{\partial \mathbb{T}^{(1)}_{(m)1}}{\partial p^1_m} - \mathbb{T}^{(1)}_{(r)1}C^{r(m)}_{m(1)} = \frac{\partial \mathbb{T}^{(1)}_{(m)1}}{\partial p^1_m},$$

$$\mathbb{T}^1_{i/1} \overset{def}{=} \frac{\delta \mathbb{T}^1_i}{\delta t} + \mathbb{T}^1_i \mathrm{K}^1_{11} - \mathbb{T}^1_r A^r_{i1} = \frac{\delta \mathbb{T}^1_i}{\delta t} + \mathbb{T}^1_i \mathrm{K}^1_{11},$$

$$\mathbb{T}^m_{i|m} \overset{def}{=} \frac{\delta \mathbb{T}^m_i}{\delta x^m} + \mathbb{T}^r_i H^m_{rm} - \mathbb{T}^m_r H^r_{im} = \mathsf{E}^m_{i|m} := \frac{\delta \mathsf{E}^m_i}{\delta x^m} + 4\mathsf{E}^m_i\sigma_m - 4\mathsf{E}^i_i\sigma_i,$$

$$\mathbb{T}^{(1)}_{(m)i}|^{(m)}_{(1)} \overset{def}{=} \frac{\partial \mathbb{T}^{(1)}_{(m)i}}{\partial p^1_m} - \mathbb{T}^{(1)}_{(r)i} C^{r(m)}_{m(1)} - \mathbb{T}^{(1)}_{(m)r} C^{r(m)}_{i(1)},$$

$$\mathbb{T}^{1(i)}_{\ \ (1)/1} \overset{def}{=} \frac{\delta \mathbb{T}^{1(i)}_{\ \ (1)}}{\delta t} + \mathbb{T}^{1(r)}_{\ \ (1)} A^i_{r1}, \quad \mathbb{T}^{m(i)}_{\ \ (1)|m} \overset{def}{=} \frac{\delta \mathbb{T}^{m(i)}_{\ \ (1)}}{\delta x^m} + 4\mathbb{T}^{m(i)}_{\ \ (1)}\sigma_m + 4\mathbb{T}^{i(i)}_{\ \ (1)}\sigma_i,$$

$$\mathbb{T}^{(1)(i)}_{(m)(1)}|^{(m)}_{(1)} \overset{def}{=} \frac{\partial \mathbb{T}^{(1)(i)}_{(m)(1)}}{\partial p^1_m} - \mathbb{T}^{(1)(i)}_{(r)(1)} C^{r(m)}_{m(1)} + \mathbb{T}^{(1)(r)}_{(m)(1)} C^{i(m)}_{r(1)}.$$

Proof. The local Einstein equations (4.4), together with some direct computations, lead us to what we were looking for. ■

4.2 Momentum electromagnetic-like geometrical model

In the paper [10], a geometrical theory for an electromagnetism depending on momenta was also created, using only a given Hamiltonian function H on the momentum phase space $J^{1*}(\mathbb{R}, M^4)$. In the background of the jet momentum Hamiltonian geometry from this paper, we work with the *electromagnetic distinguished* 2*-form* (sum by i and j)

$$\mathbb{F} = F^{(i)}_{(1)j} \delta p^1_i \wedge dx^j,$$

where (sum by r and m)

$$F^{(i)}_{(1)j} = \frac{h^{11}}{2}\left[\overset{*}{g}{}^{jr} \underset{2}{N}{}^{(1)}_{(r)i} - \overset{*}{g}{}^{ir} \underset{2}{N}{}^{(1)}_{(r)j} + \left(\overset{*}{g}{}^{jr} H^m_{ri} - \overset{*}{g}{}^{ir} H^m_{rj}\right) p^1_m\right].$$

The above electromagnetic components depending on momenta are characterized by some natural *geometrical Maxwell-like equations* (for more details, see Oană and Neagu [9], [10]).

By a direct calculation, we see that the (t, x)-conformal deformed Berwald-Moór Hamiltonian metric of order four (1.3) produces null momentum electromagnetic components: $F^{(i)}_{(1)j} = 0$. Consequently, our dual jet (t, x)-conformal deformed Berwald-Moór Hamiltonian geometrical electromagnetic theory is trivial one. Probably, this fact suggests that the dual jet (t, x)-conformal deformed Berwald-Moór Hamiltonian structure (1.3) has rather gravitational connotations than electromagnetic ones on the momentum phase space $J^{1*}(\mathbb{R}, M^4)$.

Acknowledgements. The authors of this paper thank to Professors Gh. Atanasiu and D.G. Pavlov for their encouragements and the useful discussions on the Berwald-Moór research topic.

References

[1] Gh. Atanasiu, V. Balan, M. Neagu, *The Pavlov's* 4*-polyform of momenta* $K(p) = \sqrt[4]{p_1 p_2 p_3 p_4}$ *and its applications in Hamilton geometry* (in Russian),

Hypercomplex Numbers in Geometry and Physics **2** (**4**), vol. **2** (2005), 134-139.

[2] Gh. Atanasiu, M. Neagu, *Canonical nonlinear connections in the multi-time Hamilton geometry*, Balkan Journal of Geometry and Its Applications, vol. **14**, no. **2** (2009), 1-12.

[3] Gh. Atanasiu, M. Neagu, *Distinguished torsion, curvature and deflection tensors in the multi-time Hamilton geometry*, Electronic Journal "Differential Geometry-Dynamical Systems", vol. **11** (2009), 20-40.

[4] L. Berwald, *Über Finslersche und Cartansche Geometrie II*, Compositio Math. **7** (1940), 141-176.

[5] G.I. Garas'ko, *The extension of the conformal transformation concept* (in Russian), Hypercomplex Numbers in Geometry and Physics **1** (**3**), vol. **2** (2005), 16-25.

[6] R. Miron, D. Hrimiuc, H. Shimada, S.V. Sabău, *The Geometry of Hamilton and Lagrange Spaces*, Kluwer Academic Publishers, 2001.

[7] A. Moór, *Über die Dualität von Finslerschen und Cartanschen Räumen*, Acta Math. **88** (1952), 347-370.

[8] M. Neagu, *Jet Finsler-like geometry for the x-dependent conformal deformation of a one-parameter family of Berwald-Moór metrics of order four*, Bulletin of the Transilvania University of Braşov, Series **III**: Mathematics, Informatics, Physics, vol. **4** (**53**), no. **2** (2011), 31-42.

[9] A. Oană, M. Neagu, *A distinguished Riemannian geometrization for quadratic Hamiltonians of polymomenta,* arXiv:1112.5442v1 [math.DG] (2011).

[10] A. Oană, M. Neagu, *From quadratic Hamiltonians of polymomenta to abstract geometrical Maxwell-like and Einstein-like equations*, arXiv:1202.4477v1 [math-ph] (2012).

[11] D.G. Pavlov, *Four-dimensional time*, Hypercomplex Numbers in Geometry and Physics **1** (**1**), vol. **1** (2004), 31-39.

[12] D.G. Pavlov, S.S. Kokarev, *Conformal gauges of the Berwald-Moór geometry and their induced non-linear symmetries* (in Russian), Hypercomplex Numbers in Geometry and Physics **2** (**10**), vol. **5** (2008), 3-14.

[13] P.K. Rashevsky, *Polymetric Geometry* (in Russian), In: "Proc. Sem. on Vector and Tensor Analysis and Applications in Geometry, Mechanics and Physics" (Ed. V.F. Kagan), vol. **5**, M-L, OGIZ, 1941.

Alexandru OANĂ and Mircea NEAGU
University Transilvania of Braşov, Department of Mathematics and Informatics, Blvd. Iuliu Maniu, no. 50, Braşov 500091, Romania.
E-mails: alexandru.oana@unitbv.ro, mircea.neagu@unitbv.ro